

2. The oligomeric compound of claim 27 wherein said J_1 is =O or =S and said J_2 is OH.
3. The oligomeric compound of claim 27 wherein said J_1 is =O, said J_2 is N (Y_0) T_0 and at least two of said N (Y_0) T_0 are the same.
4. The oligomeric compound of claim 27 wherein said J_1 is =O, said J_2 is N (Y_0) T_0 and wherein at least two of said N (Y_0) T_0 are different.
5. The oligomeric compound of claim 27 wherein each of said R_1 are the same.
6. The oligomeric compound of claim 27 wherein at least two of said R_1 are different.
7. The oligomeric compound of claim 27 wherein each of said aminodiol monomer subunits are the same.
8. The oligomeric compound of claim 27 wherein at least two of said aminodiol monomer subunits are different.
9. 2. The library of claim 28 wherein said J_1 is =O or =S and said J_2 is OH.
11. The library of claim 28 wherein said J_1 is =O, said J_2 is N (Y_0) T_0 and at least two of said N (Y_0) T_0 are the same.
12. The library of claim 28 wherein said J_1 is =O, said J_2 is N (Y_0) T_0 and at least two of said N (Y_0) T_0 are different.
13. The library of claim 28 wherein each of said R_1 is the same.
14. The library of claim 28 wherein at least two of said R_1 are different.
16. 3. The process of claim 29 wherein said step (g) is conducted after said step (b).

17. The process of claim 29 wherein said step (g) is conducted prior to step (d) for the addition of at least one monomeric subunit to said oligomeric compound.

18. The process of claim 29 wherein said step (g) is conducted prior to each iteration of said step (d).

19. The process of claim 29 wherein said step (g) is conducted only after at least one iteration of said step (e).

20. The process of claim 29 wherein said step (g) is conducted after said step (f) for the addition of at least one monomeric subunit to said oligomeric compound.

22. The process of claim 30 wherein said step (g) is conducted after said step (b).


23. The process of claim 30 wherein said step (g) is conducted prior to step (e) for the addition of at least one monomeric subunit to each of said oligomeric compounds.

24. The process of claim 30 wherein said step (g) is conducted prior to each iteration of said step (e).

25. The process of claim 30 wherein said step (g) is conducted only after at least one iteration of said step (e).

26. The process of claim 30 wherein said step (g) is conducted after said step (f) for the addition of at least one monomeric subunit to each of said oligomeric compounds.

--27 (New) An oligomeric compound comprising a plurality of aminodiols monomer subunits joined by linking groups, wherein each of said aminodiols monomer subunits has one of the structures I, II, III, IV, V or VI:



wherein:

each x is, independently, 0 to 5;

R₁ is -T-L or a base labile protecting group;

T is a single bond, a methylene group or a group having formula:



wherein:

R₁₀ is =O, =S, or =NR₁₁;

R₅ and E, independently, are a single bond, CH=CH, C≡C, O, S, NR₁₁, or C₆-C₁₄

aryl;

each R_6 , R_7 , R_8 , R_9 , R_{11} , R_{12} and R_{14} are, independently, H, alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;

m and n, independently, are 0 to 5;


p is 0 or 1;

q is 1 to about 10;

L is H, substituted or unsubstituted C_2 - C_{10} alkyl, substituted or unsubstituted C_2 - C_{10} alkenyl, substituted or unsubstituted C_2 - C_{10} alkynyl, substituted or unsubstituted C_4 - C_7 carbocyclic alkyl, substituted or unsubstituted C_4 - C_7 carbocyclic alkenyl, substituted or unsubstituted C_4 - C_7 carbocyclic alkynyl, substituted or unsubstituted C_6 - C_{14} aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), amide ($CONR_{12}$), amidine ($C(=NH)NR_{12}R_{13}$), guanidine ($NHC(=NH)NR_{12}R_{13}$), glutamyl ($R_{12}OOCCH(NR_{12}R_{13})(CH_2)_2C(=O)$), nitrate (ONO_2), nitro (NO_2), nitrile (CN), trifluoromethyl (CF_3), trifluoromethoxy (OCF_3), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH_2), azido (N_3), hydrazino ($NHNH_2$), hydroxylamino (ONH_2), sulfoxide (SO), sulfone (SO_2), sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and alkynyl groups;

R_2 is hydrogen or C_1-C_{10} alkyl.

R_3 and R_4 are independently hydrogen, an acid labile hydroxyl protecting group, a linking group or a conjugate group, wherein said linking group has the formula:



wherein:

J_1 is $=O$ or $=S$;

J_2 is OH or $N(Y_0)T_0$;

Y_0 is H or $(Q_2)_j-Z_2$;

T_0 is $(Q_1)_k-Z_1$, or together Y_0 and T_0 are joined in a nitrogen heterocycle;







Q_1 and Q_2 independently are C_2-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_4-C_7 carbocyclo alkyl, C_4-C_7 carbocyclo alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7-C_{14} aralkyl;

j and k independently are 0 or 1;

Z_1 and Z_2 independently are H , C_1-C_2 alkyl, C_2-C_{20} alkenyl, C_2-C_{20} alkynyl, C_6-C_{14} aryl, C_7-C_{15} aralkyl, halogen, $CH=O$, OR_{12} , SR_{12} , $NR_{12}R_{13}$, $C(=NH)NR_{12}R_{13}$, $CH(NR_{12}R_{13})$, $NHC(=NH)NR_{12}R_{13}$, $CH(NH_2)C(=O)OH$, $C(=O)NR_{12}R_{13}$, $C(=O)OR_{12}$, a metal coordination group, a reporter group, a nitrogen-containing heterocycle, a purine, a pyrimidine, a phosphate group, a polyether group, or a polyethylene glycol group; and

provided that at least one of said aminodiol monomer subunits in said oligomeric compound does not have structure III.

28 (New) A library of oligomers, each of said oligomers comprising a plurality of aminodiol monomer subunits joined by linking groups, said aminodiol monomer subunits, each of said subunits having structure I, II, III, IV, V or VI.



wherein:

each x is, independently, 0 to 5;

R₁ is -T-L or a base labile protecting group.

T is a single bond, a methylene group or a group having formula:



wherein:

R_{10} is $=\overset{\text{N}}{\underset{\text{O}}{\text{C}}}$, $=\text{S}$, or $=\text{NR}_{11}$;

R_5 and E, independently, are a single bond, $\text{CH}=\text{CH}$, $\text{C}\equiv\text{C}$, O, S, NR_{11} , or $\text{C}_6\text{-C}_{14}$ aryl;

each R_6 , R_7 , R_8 , R_9 , R_{11} , R_{12} and R_{13} are, independently, H, alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;

m and n, independently, are 0 to 5;

p is 0 or 1;


q is 1 to about 10;

L is H, substituted or unsubstituted $\text{C}_2\text{-C}_{10}$ alkyl, substituted or unsubstituted $\text{C}_2\text{-C}_{10}$ alkenyl, substituted or unsubstituted $\text{C}_2\text{-C}_{10}$ alkynyl, substituted or unsubstituted $\text{C}_4\text{-C}_7$ carbocyclic alkyl, substituted or unsubstituted $\text{C}_4\text{-C}_7$ carbocyclic alkenyl, substituted or unsubstituted $\text{C}_4\text{-C}_7$ carbocyclic alkynyl, substituted or unsubstituted $\text{C}_6\text{-C}_{14}$ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto ($\text{C}=\text{O}$), carboxyl (COOH), amide (CONR_{12}), amidine ($\text{C}(=\text{NH})\text{NR}_{12}\text{R}_{13}$), guanidine ($\text{NHC}(=\text{NH})\text{NR}_{12}\text{R}_{13}$), glutamyl ($\text{R}_{12}\text{OOCCH}(\text{NR}_{12}\text{R}_{13})(\text{CH}_2)_2\text{C}(=\text{O})$), nitrate (ONO_2), nitro (NO_2), nitrile (CN), trifluoromethyl (CF_3), trifluoromethoxy (OCF_3), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH_2), azido (N_3), hydrazino (NHNH_2), hydroxylamino (ONH_2), sulfoxide (SO), sulfone (SO_2), sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side

chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and alkynyl groups;

R_2 is hydrogen or C_1 - C_{10} alkyl;

R_3 and R_4 are independently hydrogen, an acid labile hydroxyl protecting group, a linking group or a conjugate group, wherein said linking group has the formula:



wherein:

J_1 is =O or =S;

J_2 is OH or $N(Y_0)T_0$;

Y_0 is H or $(Q_2)_j-Z_2$;

T_0 is $(Q_1)_k-Z_1$, or together Y_0 and T_0 are joined in a nitrogen heterocycle;

Q_1 and Q_2 independently are C_2 - C_{10} alkyl, C_2 - C_{10} alkenyl, C_2 - C_{10} alkynyl, C_4 - C_7 carbocyclo alkyl, C_4 - C_7 carbocyclo alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7 - C_{14} aralkyl;

j and k independently are 0 or 1;







Z_1 and Z_2 independently are H, C_1 - C_2 alkyl, C_2 - C_{20} alkenyl, C_2 - C_{20} alkynyl, C_6 - C_{14} aryl, C_7 - C_{15} aralkyl, halogen, $CH=O$, OR_{12} , SR_{12} , $NR_{12}R_{13}$, $C(=NH)NR_{12}R_{13}$, $CH(NR_{12}R_{13})$, $NHC(=NH)NR_{12}R_{13}$, $CH(NH_2)C(=O)OH$, $C(=O)NR_{12}R_{13}$, $C(=O)OR_{12}$, a metal coordination

group, a reporter group, a nitrogen-containing heterocycle, a purine, a pyrimidine, a phosphate group, a polyether group, or a polyethylene glycol group; and

provided that at least one of said aminodiols monomer subunits in each oligomeric compound of said library does not have structure III.

29. (New) A method for preparing an oligomer comprising:

(a) selecting an aminodiols monomer subunit having the structure I, II, III, IV, V, or VI:



wherein:

each x is, independently, 0 to 5;

R₁ is a base labile amino protecting group;

R₂ is hydrogen or C₁-C₁₀ alkyl.

one of R_1 or R_4 is hydrogen or an activated phosphite group and the other of R_1 or R_4 is an acid labile hydroxyl protecting group.


(b) attaching said aminodiol monomer subunit to a solid support to form a solid support bound aminodiol monomer subunit.

(c) treating said acid labile hydroxyl protecting group with a dilute acid to form a free hydroxyl group.

(d) reacting said free hydroxyl group with a further aminodiol monomer subunit having structure I, II, III, IV, V, or VI thereby forming an oligomeric compound bound to said solid support, said oligomeric compound containing a phosphite linkage:

(e) optionally iteratively repeating steps (c) and (d) to increase the length of the oligomeric compound bound to said solid support.

(f) optionally, prior to step (e) or after step (d) oxidizing said phosphite linkage to form a phosphate linking group wherein said linking groups are selected having formula:



wherein:

J_1 is =O or =S;

J_2 is OH or $N(Y_0)T_0$;

Y_0 is H or $(Q_2)_j-Z_2$;

T_0 is $(Q_1)_k-Z_1$, or together Y_0 and T_0 are joined in a nitrogen heterocycle;

Q_1 and Q_2 independently are C_2-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_4-C_7 carbocyclo alkyl, C_4-C_7 carbocyclo alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7-C_{14} aralkyl;

j and k independently are 0 or 1;

Z_1 and Z_2 independently are H, C_1-C_2 alkyl, C_2-C_{20} alkenyl, C_2-C_{20} alkynyl, C_6-C_{14} aryl, C_7-C_{15} aralkyl, halogen, $CH=O$, OR_{12} , SR_{12} , $NR_{12}R_{13}$, $C(=NH)NR_{12}R_{13}$, $CH(NR_{12}R_{13})$, $NHC(=NH)NR_{12}R_{13}$, $CH(NH_2)C(=O)OH$, $C(=O)NR_{12}R_{13}$, $C(=O)OR_{12}$, a metal coordination group, a reporter group, a nitrogen-containing heterocycle, a purine, a pyrimidine, a phosphate group, a polyether group, or a polyethylene glycol group;

(g) prior to step (e) or after step (f) contacting said solid support bound aminodiol monomer subunit or said support bound oligomeric compound with a base to remove said base labile amino protecting group to form the solid support bound aminodiol monomer subunit or support bound oligomeric compound having a free amine, and derivatizing said free amine with a group of the formula:



wherein:

T is a single bond, a methylene group or a group having formula:



where:

R_{10} is $=O$, $=S$, or $=NR_{11}$;

R_5 and E, independently, are a single bond, $\text{CH}=\text{CH}$, $\text{C}\equiv\text{C}$, O, S, NR_{11} , or $\text{C}_6\text{-C}_{14}$ aryl.

each R_6 , R_7 , R_8 , R_9 , R_{11} , R_{12} and R_{13} are, independently, H, alkyl or haloalkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;

m and n, independently, are 0 to 5;

p is 0 or 1;

q is 1 to about 10.

L is H, substituted or unsubstituted $\text{C}_2\text{-C}_{10}$ alkyl, substituted or unsubstituted $\text{C}_2\text{-C}_{10}$ alkenyl, substituted or unsubstituted $\text{C}_2\text{-C}_{10}$ alkynyl, substituted or unsubstituted $\text{C}_4\text{-C}_7$ carbocyclic alkyl, substituted or unsubstituted $\text{C}_4\text{-C}_7$ carbocyclic alkenyl, substituted or unsubstituted $\text{C}_4\text{-C}_7$ carbocyclic alkynyl, substituted or unsubstituted $\text{C}_6\text{-C}_{14}$ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto ($\text{C}=\text{O}$), carboxyl (COOH), amide (CONR_{12}), amidine ($\text{C}(=\text{NH})\text{NR}_{12}\text{R}_{13}$), guanidine ($\text{NHC}(=\text{NH})\text{NR}_{12}\text{R}_{13}$), glutamyl ($\text{R}_{12}\text{OOCCH}(\text{NR}_{12}\text{R}_{13})(\text{CH}_2)_2\text{C}(=\text{O})$), nitrate (ONO_2), nitro (NO_2), nitrile (CN), trifluoromethyl (CF_3), trifluoromethoxy (OCF_3), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH_2), azido (N_3), hydrazino (NHNH_2), hydroxylamino (ONH_2), sulfoxide (SO), sulfone (SO_2), sulfide (S-), disulfide (S-S), silyl, a nucleosidic base, an amino acid side chain, a carbohydrate, a biopharmaceutically active moiety, or group capable of hydrogen

bonding where the substituent groups are selected from hydroxyl, amino, alkoxy, alcohol,

benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl, and alkynyl groups.







(h) optionally repeating steps (c) and (d) followed by step (g) to increase the length of the oligomeric compound bound to said solid support;

(i) treating said oligomeric compound bound to said solid support with acid to deprotect any protecting groups; and

(j) cleaving said oligomeric compound from said solid support.

30. A method for preparing a combinatorial library comprising:

(a) selecting a plurality of aminodiols monomer subunits having the structure I, II, III, IV, V, or VI:



wherein:

each x is, independently, 0 to 5.

R_1 is a base labile amino protecting group.

R_2 is hydrogen or C_1 - C_{10} alkyl.

one of R_3 or R_4 is hydrogen or an activated phosphite group and the other of R_3 or R_4 is an acid labile hydroxyl protecting group.


(b) attaching said aminodiol monomer subunits to a solid support to form a solid support bound aminodiol monomer subunits;

(c) treating said acid labile hydroxyl protecting groups with a dilute acid to form a free hydroxyl groups;

(d) reacting said free hydroxyl groups with further aminodiol monomer subunits having structure I, II, III, IV, V or VI thereby forming an oligomeric compound bound to said solid support, said oligomeric compound containing a phosphite linkage;

(e) optionally iteratively repeating steps (c) and (d) to increase the length of the oligomeric compound bound to said solid support;

(f) optionally, prior to step (c) or after step (d) oxidizing said phosphite linkage to form phosphate linking groups having formula:



wherein:

J_1 is =O or =S;

J_2 is OH or $N(Y_n)T_n$;

Y_n is H or $(Q_2)_j-Z_2$;

T_n is $(Q_1)_k-Z_1$, or together Y_n and T_n are joined in a nitrogen heterocycle;

Q_1 and Q_2 independently are C_2-C_{10} alkyl, C_2-C_{10} alkenyl, C_2-C_{10} alkynyl, C_4-C_7 carbocyclo alkyl, C_4-C_7 carbocyclo alkenyl, a heterocycle, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a polyalkyl glycol, or C_7-C_{14} aralkyl;

j and k independently are 0 or 1;

Z_1 and Z_2 independently are H, C_1-C_2 alkyl, C_2-C_{20} alkenyl, C_2-C_{20} alkynyl, C_6-C_{14} aryl, C_7-C_{15} aralkyl, halogen, $CH=O$, OR_{12} , SR_{12} , $NR_{12}R_{13}$, $C(=NH)NR_{12}R_{13}$, $CH(NR_{12}R_{13})$, $NHC(=NH)NR_{12}R_{13}$, $CH(NH_2)C(=O)OH$, $C(=O)NR_{12}R_{13}$, $C(=O)OR_{12}$, a metal coordination group, a reporter group, a nitrogen-containing heterocycle, a purine, a pyrimidine, a phosphate group, a polyether group, or a polyethylene glycol group;

(g) prior to step (e) or after step (f) contacting said solid support bound aminodiol monomer subunits or said support bound oligomeric compounds with a base to remove said base labile amino protecting groups to form the solid support bound aminodiol monomer subunits or support bound oligomeric compounds having a free amine, and derivatizing said free amine with a group of the formula

-T-L

wherein:

T is a single bond, a methylene group or a group having formula:



where:

R_{10} is =O, =S, or =NR₁₁;

R₅ and E, independently, are a single bond, CH=CH, C≡C, O, S, NR₁₁, or C₆-C₁₄

aryl;

each R₆, R₇, R₈, R₉, R₁₁, R₁₂ and R₁₃ are, independently, H, alkyl or haloalkyl

having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, alkynyl having 2 to about 10 carbon atoms, or aryl having 7 to about 14 carbon atoms;

m and n, independently, are 0 to 5;

p is 0 or 1;

q is 1 to about 10;

L is H, substituted or unsubstituted C₂-C₁₀ alkyl, substituted or unsubstituted C₂-C₁₀ alkenyl, substituted or unsubstituted C₂-C₁₀ alkynyl, substituted or unsubstituted C₄-C₇ carbocyclic alkyl, substituted or unsubstituted C₄-C₇ carbocyclic alkenyl, substituted or unsubstituted C₄-C₇ carbocyclic alkynyl, substituted or unsubstituted C₆-C₁₄ aryl, an ether having 2 to 10 carbon atoms and 1 to 4 oxygen or sulfur atoms, a nitrogen containing heterocycle, a sulfur containing heterocycle, an oxygen containing heterocycle, a metal coordination group, a conjugate group, halogen, hydroxyl (OH), thiol (SH), keto (C=O), carboxyl (COOH), amide (CONR₁₂), amidine (C(=NH)NR₁₂R₁₃), guanidine (NHC(=NH)NR₁₂R₁₃), glutamyl (R₁₂OOCCCH(NR₁₂R₁₃)(CH₂)₂C(=O)), nitrate (ONO₂), nitro (NO₂), nitrile (CN), trifluoromethyl (CF₃), trifluoromethoxy (OCF₃), O-alkyl, S-alkyl, NH-alkyl, N-dialkyl, O-aralkyl, S-aralkyl, NH-aralkyl, amino (NH₂), azido (N₃), hydrazino (NHNH₂), hydroxylamino (ONH₂), sulfoxide